

CLINICAL IMAGES**Congenital lung mass in an asymptomatic patient**

Heather Zar, Bruce McIvor, Gisella Furlan, Leon Jedeikin, Richard Pitcher

A routine 20-week antenatal ultrasound scan showed a congenital lesion of the left fetal lung, measuring approximately 25 mm x 25 mm x 30 mm. The mass showed no sonographic change through the remainder of an uneventful pregnancy. The baby was delivered by elective caesarean section at 38 weeks' gestation, with a birth weight of 2 900 g, and had no postnatal complications. A chest radiograph performed in the early neonatal period was normal, but a contrasted single-slice helical computed tomography (CT) chest scan at age 6 weeks demonstrated the small, oval, solid mass in the left lower lobe, with no associated mediastinal shift (Fig. 1). The vascular supply of the lesion could not be identified on this scan.

Over the ensuing 9 months the baby has been well, without respiratory symptoms.

A follow-up contrasted CT scan was performed at age 7 months on a 64-slice multi-detector scanner. This showed the mass to be morphologically unchanged. However, the multiplanar reconstructions possible on the multislice scanner facilitated a definitive diagnosis. The mass was shown to have a systemic arterial supply, via a 3 mm diameter vessel arising from the descending thoracic aorta, with venous drainage through 3 radicals into the hemi-azygous vein (Figs 2 - 3). These findings are consistent with extralobar pulmonary sequestration.

Professor Heather Zar is a paediatric pulmonologist at Red Cross Children's Hospital and the University of Cape Town.

Bruce McIvor is a general radiologist in private practice with Dr Morton and Partners in Cape Town. Special interests include multislice CT, in particular cardiac CT, and MRI.

Gisella Furlan is a full-time radiologist at Red Cross Children's Hospital.

Leon Jedeikin is a paediatrician in private practice at Vincent Pallotti Hospital in Cape Town and a part-time tutor and geneticist at Red Cross Children's hospital.

Richard Pitcher is Head of the Division of Paediatric Radiology at Red Cross Children's Hospital and the University of Cape Town.




Fig. 1. Axial CT scan through the lower chest, demonstrating the oval, solid mass in the left lower lobe, posteriorly (white arrow). Note also the prominent hemi-azygous vein (black arrow), running dorsal to the descending thoracic aorta (A).




Fig. 2. Reconstructed oblique image demonstrating the systemic arterial supply to the congenital pulmonary lesion. Note the supplying artery (white arrow), arising from the abdominal aorta (A) coursing upwards into the chest to supply the solid mass (M) at the lung base (L).

Discussion**Radiological advances**

This case report indicates how radiological advances have enhanced both antenatal identification of congenital pulmonary abnormalities and their postnatal diagnosis. Previously such asymptomatic pulmonary lesions may have remained

Corresponding author: R Pitcher (pitcher@iafrica.com)




Fig. 3. Reconstructed parasagittal image, demonstrating the venous drainage of the lesion. Note confluence of three draining venules (white arrow), into the hemi-azygous vein (black arrow), lying dorsal to the descending thoracic aorta (A).

undetected until the onset of complications in later life. Furthermore a specific diagnosis would have required invasive procedures in the form of angiography or biopsy.

Radiation risk from paediatric CT has received particular attention in the last 5 years,¹ prompting strategies for dose reduction that will largely preserve image quality. As with any radiological procedure, chest CT should be used for appropriate clinical indications, bearing in mind that the radiation dose is equivalent to approximately 250 standard antero-posterior chest radiographs.

Pathology and management issues

The spectrum of congenital lung lesions includes sequestration, congenital cystic adenomatoid malformation (CCAM), lobar emphysema and bronchogenic cysts.² Classically, sequestration describes a bronchopulmonary mass with either no connection, or an abnormal connection to the bronchial tree. It typically has an anomalous systemic arterial supply,³ and may be intralobar (within the pleura of the involved lung) or extralobar. An intralobar sequestration usually has normal pulmonary venous drainage, while the extralobar lesion characteristically has aberrant drainage into the right atrium, vena cava or azygous systems, as in this patient.^{2,4}

Extralobar sequestrations are less common than their intralobar counterparts, have a 4:1 male to female preponderance and occur on the left side in 90% of cases.⁵ Up to 65% are associated with other congenital anomalies. Since congenital pulmonary lesions may have overlapping

features, the broader term 'congenital thoracic malformation' is recommended in preference to applying a pathological diagnosis in a clinical setting.²

Many such malformations detected at the 20-week antenatal scan regress or disappear by term.⁶ Although sequestrations are rarely symptomatic at birth,⁷ symptoms can be severe and are determined by the size and location of the lesion.^{2,3} Initially asymptomatic lesions may become symptomatic owing to the development of complications, in which case surgical removal is indicated.⁸

Therapy for asymptomatic lesions is controversial and the optimal management strategy is not clearly defined. Options include conservative management with careful long-term follow-up, surgical removal, or embolisation of the systemic feeder vessel. Reasons for surgical intervention or embolisation include the risk of recurrent infections, an increase in the arteriovenous shunt, pressure effect on adjacent normal lung, airway compression and malignant degeneration.⁸⁻¹⁰ The latter has been reported in congenital malformations, particularly CCAMs.^{9,10}

Some experts advocate resection of all lesions irrespective of symptoms because of the potential for complications, while others have promoted conservative management owing to the lack of data on the long-term natural history of asymptomatic lesions and the possibility of their spontaneous regression.⁶ The optimal timing of surgery is also unclear.¹¹

The authors record their thanks to the parents for permission to publish.

- Brenner DJ, Elliston CD, Hall EJ, Berdon WE. Estimated risks of radiation induced fatal cancer from paediatric CT. *Am J Roentgenol* 2001; **176**: 289-296.
- Bush A. Congenital lung disease: a plea for clear thinking and clear nomenclature. *Pediatr Pulmonol* 2001; **32**: 328-337.
- Horak E, Bodner J, Gassner I, et al. Congenital cystic lung disease: diagnostic and therapeutic considerations. *Clin Pediatr (Phila)* 2003; **42**: 251-261.
- Clements BS, Warner JO. Pulmonary sequestration and related congenital bronchopulmonary-vascular malformations: nomenclature and classification based on anatomical and embryological considerations. *Thorax* 1987; **42**: 401-408.
- Bercoff T, Madrid C, Novo S, Gutierrez J, Arjonilla A, Gomez-Leon N. Congenital anomalies of the tracheobronchial tree, lung and mediastinum: embryology, radiology and pathology. *Radiographics* 2004; **24**(1): e 17. Epub 2003, Nov. 10. Review.
- Lacy DE, Shaw NJ, Pilling DW, Walkinshaw S. Outcome of congenital lung abnormalities detected antenatally. *Acta Paediatr* 1999; **88**: 454-458.
- Piccione W jun., Burt ME. Pulmonary sequestration in the neonate. *Chest* 1990; **97**: 244-246.
- Al-Bassam A, Al-Rabeeh A, Al-Nassar S, et al. Congenital cystic disease of the lung in infants and children (experience with 57 cases). *Eur J Pediatr Surg* 1999; **9**: 364-368.
- Papagiannopoulos K, Hughes S, Nicholson AG, Goldstraw P. Cystic lung lesions in the pediatric and adult population: surgical experience at the Brompton Hospital. *Ann Thorac Surg* 2002; **73**: 1594-1598.
- MacSweeney F, Papagiannopoulos K, Goldstraw P, Sheppard MN, Corrin B, Nicholson AG. An assessment of the expanded classification of congenital cystic adenomatoid malformations and their relationship to malignant transformation. *Am J Surg Pathol* 2003; **27**: 1139-1146.
- Laberge JM, Puligandla P, Flageole H. Asymptomatic congenital lung malformations. *Semin Pediatr Surg* 2005; **14**(1): 16-33.