

# Mapping evaporative water loss in desert passerines reveals an expanding threat of lethal dehydration

Thomas P. Albright<sup>a,b,1</sup>, Denis Mutiibwa<sup>a,c</sup>, Alexander. R. Gerson<sup>d,e</sup>, Eric Krabbe Smith<sup>e</sup>, William A. Talbot<sup>e</sup>, Jacqueline J. O'Neill<sup>e</sup>, Andrew E. McKechnie<sup>f</sup>, and Blair O. Wolf<sup>e,1</sup>

<sup>a</sup>Department of Geography, University of Nevada, Reno, NV 89557; <sup>b</sup>Program in Ecology, Evolution, and Conservation Biology, University of Nevada, Reno, NV 89557; <sup>c</sup>Beltsville Agricultural Research Center, US Department of Agriculture, Beltsville Agricultural Research Center–West, Beltsville, MD 20705; <sup>d</sup>Department of Biology, University of Massachusetts, Amherst, MA 01003; <sup>e</sup>Biology Department, University of New Mexico, Albuquerque, NM 87131-0001; and <sup>f</sup>DST-NRF Centre of Excellence at the Percy FitzPatrick Institute, Department of Zoology and Entomology, University of Pretoria, Hatfield 0028, South Africa

Edited by Scott R. McWilliams, University of Rhode Island, Kingston, RI, and accepted by Editorial Board Member Gregory P. Asner December 21, 2016 (received for review August 16, 2016)

**Extreme high environmental temperatures produce a variety of consequences for wildlife, including mass die-offs. Heat waves are increasing in frequency, intensity, and extent, and are projected to increase further under climate change. However, the spatial and temporal dynamics of die-off risk are poorly understood. Here, we examine the effects of heat waves on evaporative water loss (EWL) and survival in five desert passerine birds across the southwestern United States using a combination of physiological data, mechanistically informed models, and hourly geospatial temperature data. We ask how rates of EWL vary with temperature across species; how frequently, over what areas, and how rapidly lethal dehydration occurs; how EWL and die-off risk vary with body mass; and how die-off risk is affected by climate warming. We find that smaller-bodied passerines are subject to higher rates of mass-specific EWL than larger-bodied counterparts and thus encounter potentially lethal conditions much more frequently, over shorter daily intervals, and over larger geographic areas. Warming by 4 °C greatly expands the extent, frequency, and intensity of dehydration risk, and introduces new threats for larger passerine birds, particularly those with limited geographic ranges. Our models reveal that increasing air temperatures and heat wave occurrence will potentially have important impacts on the water balance, daily activity, and geographic distribution of arid-zone birds. Impacts may be exacerbated by chronic effects and interactions with other environmental changes. This work underscores the importance of acute risks of high temperatures, particularly for small-bodied species, and suggests conservation of thermal refugia and water sources.**

avian ecology | physiological ecology | climate change | heat waves | water balance

Extreme weather events are increasingly seen as an important factor in ecology and conservation, with consequential effects on individuals, populations, communities, and ecosystems (1–3). Recent data indicate an increase in the incidence of heat waves and extreme high temperatures (4, 5). Despite difficulties in quantifying trends in mass mortality events, heat waves are known to have caused a number of large-scale die-offs among birds, pteropodid bats, and other taxa in recent years (6, 7). Moreover, current (8) and projected (9) increases in the frequency, duration, and severity of heat waves are likely to make these mortality events more common as the century progresses (10).

Birds may be particularly susceptible to heat waves given their typically diurnal activity periods, small size, and high mass-specific rates of metabolism and water loss. Small birds also have a very limited capacity to store vital resources such as water, and consequently must balance their water budgets over time scales of minutes to hours during hot weather (10). Constraints on water availability and heat stress are known to produce changes in behavior, reproductive success, occupancy, and mortality in birds (11). Heat-related mortality can arise from two categories of physiological constraints: (i) limits to the balancing of water budgets and

maintenance of adequate hydration under conditions requiring evaporative cooling (dehydration), and (ii) limits to rates of heat dissipation and the maintenance of body temperature ( $T_b$ ) below lethal limits (hyperthermia). When environmental temperatures exceed a bird's normothermic  $T_b$ , evaporative heat loss becomes the only avenue of heat dissipation available to offset increasing environmental and internal heat loads. Rates of evaporative water loss (EWL) increase markedly when environmental temperatures are above  $T_b$  and can exceed 5% of body mass ( $M_b$ ) per hour, leading to rapid and acute losses of water (12); this creates a fundamental tradeoff between the dangers of dehydration and of hyperthermia. Given these challenges, birds typically curtail activity during the hottest periods of the day and seek relatively cool, shaded microsites to minimize heat loads and EWL (13).

Body mass is one of the most important factors affecting bird–environment interactions, with smaller species showing higher mass-specific rates of EWL and hence greater relative water demands. In a modeling study, McKechnie and Wolf (10) showed that warming temperatures in hot deserts could be expected to produce large fractional increases in EWL rates, and that future extreme heat events could result in very short survival times for smaller birds (<2 h for birds <10 g). However, this study was limited to physiological data obtained from the literature, and models were implemented for only two sites. Furthermore, the authors did not address the effects of geographic variation in

## Significance

Using measured rates of evaporative water loss, hourly gridded weather data, a 4 °C warming scenario, and physiological models, we show that songbirds in the deserts of the southwestern United States are increasingly susceptible to death from dehydration on hot days. Smaller birds lose water at a proportionally higher rate, and are hence more vulnerable than larger birds to lethal dehydration arising from greater evaporative cooling demands. Our analysis indicates that, by the end of the present century, exposure to potentially lethal conditions could at least quadruple for smaller species. The increasing extent, frequency, and intensity of dehydrating conditions under a warming climate may alter daily activity patterns, geographic range limits, and the conservation status of affected birds.


Author contributions: T.P.A., A.E.M., and B.O.W. designed research; D.M., A.R.G., E.K.S., W.A.T., J.J.O., A.E.M., and B.O.W. performed research; D.M. contributed new analytic tools; T.P.A., D.M., A.R.G., and B.O.W. analyzed data; and T.P.A., D.M., A.R.G., A.E.M., and B.O.W. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission. S.R.M. is a Guest Editor invited by the Editorial Board.

<sup>1</sup>To whom correspondence may be addressed. Email: talbright@unr.edu or wolf@unm.edu.

This article contains supporting information online at [www.pnas.org/lookup/suppl/doi:10.1073/pnas.1613625114/-DCSupplemental](http://www.pnas.org/lookup/suppl/doi:10.1073/pnas.1613625114/-DCSupplemental).



**Fig. 1.** Evaporative water loss (EWL) as a percentage of body mass ( $M_b$ ) for five passerine bird species across a range of elevated chamber  $T_{air}$ . Mean body masses ( $M_b$ ) are indicated and a linear fit line ( $P < 0.001$ ) has been added for  $T_{air}$  above  $T_{uc}$  of 40 °C. A small amount of jitter (10% minimum distance between adjacent unique values) has been added to minimize overplotting.


temperature across major portions of species' ranges, thus excluding a dimension critical for conservation planning.

Here, we examine the effects of heat waves on rates of EWL and survival in desert-dwelling birds in the United States using a combination of physiological data, mechanistically informed models, and hourly geospatial temperature data. We use measurements of EWL as a function of  $T_{air}$  above 40 °C (the approximate diurnal  $T_b$  of desert passerines) for five passerine birds common to the desert southwest. We model EWL and map the occurrence of current (1980–2013) conditions of lethal dehydration, which we define as cumulative EWL exceeding 15% of  $M_b$ . Given daily activity patterns and temperature profiles, we consider survival times of  $\leq 5$  h to pose an ecologically important (moderate) risk of lethal dehydration, and base most of our spatial modeling on this scenario. In addition, we characterize the occurrence of survival times  $\leq 3$  h (high risk). We then explore the implications of a 4 °C warming scenario, consistent with climate projections for this region in 2070–2100 (9, 14). Using


these five passerines, we ask (i) How do rates of EWL vary as a function of  $M_b$  across a range of  $T_{air}$ ? (ii) How frequently, in what areas, and over what time scales do conditions associated with lethal dehydration currently occur? (iii) How will a warmer climate affect the frequency and spatial extent of conditions under which lethal dehydration occurs?

## Results

**Evaporative Water Loss Rates as a Function of Body Mass and  $T_{air}$ .** At  $T_{air} < 40$  °C, rates of EWL were universally well below  $1 \text{ g}\cdot\text{h}^{-1}$ , with larger species having slightly higher rates than smaller species. However, at  $T_{air} > 40$  °C, rates of total EWL increased markedly in a linear fashion. In the largest species, the Curve-billed Thrasher (71 g), EWL exceeded  $4 \text{ g}\cdot\text{h}^{-1}$  at 50 °C. However, small species evaporated water at proportionally higher rates than larger species (Fig. 1). For example, at 50 °C, Lesser Goldfinch (9.7 g) and House Finch (18.0 g) evaporated 8–9% of  $M_b\cdot\text{h}^{-1}$ , whereas the larger Abert's Towhee (41.8 g)



**Fig. 2.** Current (1980–2012) and future (2070–2100) number of days between April 1 and September 30 with survival time of  $\leq 5$  h across the range (orange lines) of three passerine species in the southwestern United States, arranged in order of increasing  $M_b$ . Maps of additional species and for  $\leq 3$ -h dehydration risk are found in Fig. S1.



**Fig. 3.** Percentage area of species ranges in the Southwest study region subject to conditions associated with lethal dehydration under current (*Upper*) and future 4 °C warmer (*Lower*) climates for moderate (survival time  $\leq 5$  h; *Left*) and high (survival time  $\leq 3$  h; *Right*) dehydration risk. LEGO, Lesser Goldfinch; HOFI, House Finch; CAWR, Cactus Wren; ABTO, Abert's Towhee; CBTH, Curve-billed Thrasher. The bottom of box, heavy central line, and top of box indicate areas experiencing  $\geq 10$  d $\cdot$ y $^{-1}$ ,  $\geq 5$  d $\cdot$ y $^{-1}$ , and  $\geq 1$  d $\cdot$ y $^{-1}$ , respectively. The width of each box is proportional to the range area of species within the conterminous United States, and abscissa position is determined by body mass.

and Curve-billed Thrasher evaporated  $\sim 7\%$  and  $5\%$  of  $M_b \cdot h^{-1}$ , respectively.

#### Spatiotemporal Dynamics of Lethal Dehydration: Current Conditions.


Maps of weather conditions associated with lethal dehydration reveal variable spatial patterns among our suite of passerine birds (Fig. 2 and Fig. S1). First, small passerines experienced lethal dehydration risk much more frequently than larger species, often with more than 50 d $\cdot$ y $^{-1}$  over large portions of their range. For instance, at Yuma, AZ (32° 42' N, 114° 37' W), Lesser Goldfinch is modeled to experience an average 58 d $\cdot$ y $^{-1}$  with survival time of  $\leq 5$  h, whereas the larger Curve-billed Thrasher only experiences 4 d $\cdot$ y $^{-1}$  of comparable risk. Second, conditions associated with high dehydration risk (i.e., survival times  $\leq 3$  h) are presently limited to only the smallest species considered, with Lesser Goldfinch experiencing only a few days per year in limited low-elevation portions of their range (Fig. 1). Among our focal species, modeled survival times are currently never  $\leq 3$  h for Abert's Towhee or Curve-billed Thrasher.

Spatially, large portions of the US ranges of these species are subject to at least occasional risk of lethal dehydration (Fig. 2). Considering the normal conterminous US breeding ranges, the percentage of area experiencing an average of at least 1 d $\cdot$ y $^{-1}$  of lethal dehydration risk ranged from 42% for Lesser Goldfinch to 20% for Abert's Towhee (Fig. 3). However, areas in which these species currently experience more than 10 d $\cdot$ y $^{-1}$  of these conditions were much more restricted. The occurrence of high lethal dehydration risk ( $\leq 3$  h) was similarly limited to parts of the Sonoran, Colorado, and Mojave deserts of western Arizona, southern Nevada, and southeastern California in smaller species, with these conditions not occurring at all in average years for larger species.

**Climate Warming Effects on Spatiotemporal Dynamics of Lethal Dehydration.** Warming of 4 °C greatly expands the extent, frequency, and intensity of dehydration risk for desert passerines in the United States. All species modeled will be subjected to increased risk of lethal dehydration over large parts of the southwestern United States (Fig. 2). Across the entire study region, there will be a large increase in the area of a species' range experiencing at least 1 d $\cdot$ y $^{-1}$  with moderate ( $< 5$  h) dehydration risk; this is particularly true for the two larger-bodied, range-restricted birds, the Cactus Wren and Curve-billed Thrasher, with  $> 80\%$  of their ranges affected. Increases in the frequency of extreme dehydration risk are perhaps more dramatic, especially for the smaller species (Fig. 2). In the southwest desert ecoregion, the average number of days per year with lethal dehydration risk increases from 13 to 44 d for Lesser Goldfinch, from 2 to 9 d for Cactus Wren, and from  $< 1$  to 4 d for Curve-billed Thrasher (Fig. 4). Days with high lethal dehydration risk will remain rare for the largest species, but for smaller species, such days become far more frequent. For Lesser Goldfinches in the southwestern deserts, high risk of lethal dehydration increases from 7 to 25 d $\cdot$ y $^{-1}$  (Fig. 4). In some portions of their range, particularly the Sonoran Desert of southwest Arizona and extreme southeastern California, the Lesser Goldfinch will experience survival times  $\leq 3$  h for virtually the entire summer and portions of spring and fall (Fig. S1).

#### Discussion

Our models reveal that increasing  $T_{air}$  and the concomitant increase in the frequency and severity of heat waves will have potentially important impacts on the water balance, daily activity, and geographic distribution of birds in deserts of the southwestern United States. Even modest increases in maximum  $T_{air}$  during heat waves result in greatly increased demands for evaporative cooling. Increases in rates of EWL are strongly dependent on  $M_b$ , with smaller species showing higher mass-specific rates. Consequently, smaller species will be subject to much greater dehydration risk than larger species. Because activity is reduced at high  $T_{air}$ , the impact of higher EWL rates will be exacerbated by reduced foraging and hence water intake. Thus, the selection of relatively cool microhabitats may be expected to become increasingly important. Finally, diet will also play an important role in determining future avian distributions, with granivorous birds challenged by their relatively dry diet and need for scarce, unpredictable free water resources. In the following paragraphs, we discuss each of these topics and how their integrative effects under conditions of rapid warming may affect the future distributions of arid-zone passerines.



**Fig. 4.** Average days per year with 5-h (*Left*) and 3-h (*Right*) lethal dehydration intensities in current and future climates. Averages are calculated over each species' range within the US hot desert ecoregion, comprising the Mojave, Sonoran, and Chihuahuan deserts (ecoregion data from ref. 39).

**Scaling of Rates of Evaporation and Time to Lethal Dehydration.** The effects of  $M_b$  on resting rates of EWL at moderate  $T_{air}$  are well described in birds (15, 16). Less well understood is how EWL and upper limits to evaporative cooling capacity vary with  $T_{air}$  above normothermic  $T_b$  (10, 17). As expected, our data confirm that smaller species lose water at higher mass-specific EWL than larger species. At  $T_a = 40^\circ\text{C}$ , for example, the Lesser Goldfinch (9.7 g) loses  $3.4\% M_b \cdot h^{-1}$ , a rate approximately twice that of the approximately sevenfold-larger Curve-billed Thrasher (71 g;  $1.8\% M_b \cdot h^{-1}$ ). At  $T_a \sim 10^\circ\text{C}$  above normothermic  $T_b$ , rates of EWL in goldfinches can exceed  $8\% M_b \cdot h^{-1}$ , and may approach  $5\% M_b \cdot h^{-1}$  in thrashers. During extreme heat events, goldfinches may reach their limits of dehydration tolerance ( $\sim 15\% M_b$ ) in less than 2 h, and even thrashers approach these same limits in as little as 3 h (Fig. 3).

In both the Lesser Goldfinch and Curve-billed Thrasher, an increase in  $T_a$  from 40 to  $44^\circ\text{C}$  results in a  $\sim 150\%$  increase in rates of EWL. However, time to death through dehydration differs greatly between these two species because of differences in mass-specific rates of EWL ( $5.4\% M_b \cdot h^{-1}$  vs.  $2.7\% M_b \cdot h^{-1}$ ; Fig. 1). These values are similar to the fractional increases in EWL required for evaporative cooling predicted by the model presented by McKechnie and Wolf (10) for 27 avian species, and provide the basis for mapping EWL with daily changes in  $T_a$  across each species' range.

**Dehydration Risk Under Current and Future Climates.** Maps of dehydration risk (Fig. 2) provide the basis for quantifying the proportion of species' ranges where lethal dehydration ( $\leq 5$  h) is likely to occur, and the frequency of occurrence ( $d \cdot y^{-1}$ ). Interactions among body mass, geographic range, and dehydration risk reveal that, under certain scenarios, the smallest, most heat-sensitive species are the least affected from a distribution and abundance standpoint. House Finches and Lesser Goldfinches provide excellent examples, both having widespread distributions that span the latitude range of the study region. Under current climate conditions, both species exceed their 5-h dehydration limits on  $1\text{--}10 d \cdot y^{-1}$  over large areas ( $\sim 1,500,000 \text{ km}^2$  and  $\sim 960,000 \text{ km}^2$ , respectively; Fig. 2 and Fig. S1). Even though these areas of risk include 25% and 42% of the total ranges for House Finches and Lesser Goldfinches, respectively, the very large ranges ( $\sim 6$  million and  $2.3$  million  $\text{km}^2$ , respectively) indicate that the majority of populations are at low risk of mortality from acute heat stress under current conditions. Under future climates, however, these large ranges may not provide adequate population buffers, given likely additional stressors. Under the  $+4^\circ\text{C}$  scenario, House Finches will experience at least  $1 d \cdot y^{-1}$  of lethal dehydration risk over  $\sim 65\%$  of the species' total range, and Lesser Goldfinches will encounter potentially lethal events over  $\sim 72\%$  of the total range.

A different scenario applies to Abert's Towhee, Cactus Wren, and Curve-billed Thrasher, all of which are arid-zone specialists with relatively small distributional ranges ( $\sim 200,000\text{--}900,000 \text{ km}^2$ ; Fig. 2 and Fig. S1). Under current conditions, the risk of lethal dehydration for these larger passerines is reduced by their modest mass-specific rates of EWL. Thus, despite being restricted to hot deserts, the fractions of their ranges (Cactus Wren 36%; Abert's Towhee 20%; Curve-billed Thrasher 31%) over which there is a risk of lethal dehydration are comparable to those of the smaller, but more widely distributed Lesser Goldfinches and House Finches; however, because their total ranges are small and mainly confined to hot deserts, a  $4^\circ\text{C}$  warming scenario produces very different consequences. Temperatures under projected warming exceed the threshold for lethal dehydration on at least a handful of days each year over the majority of each species' range (Fig. 3; Cactus Wren 83%; Abert's Towhee 58%; and the Curve-billed Thrasher 83%). Possibly of even greater concern, however, is the observation that  $4^\circ\text{C}$  warming in an already hot desert will yield long periods when chronic effects of heat and water limitation may pose problems, as discussed below.

**Expanding Risks Under Climate Change.** Our model reveals that the portion of species' ranges with risk of lethal dehydration during heat waves will dramatically expand under future climate scenarios (Fig. 3), as will the average frequency at which such conditions are encountered (Fig. 4), and this is most apparent for the Lesser Goldfinch. By 2100, this species will be exposed to  $60\text{--}120 d \cdot y^{-1}$  of potentially lethal conditions over a large region that includes the entire lower Colorado River basin, southern Arizona, the lower central valley of California and southern Texas (Fig. 2). A similar pattern emerges for House Finches, where in the central valley of California and the entire lower Colorado River basin there is a sustained risk of lethal dehydration much of the year. Under the current climate, exposure of these two species to such extreme conditions is mostly limited to the lower Colorado River basin and constrained to  $20\text{--}40 d \cdot y^{-1}$ . As granivores, these two species may be at particular risk because of their reliance on free water sources, an ecological trait that may well exclude them from large parts of their current range by the end of this century. For the larger three species, Cactus Wren, Abert's Towhee, and Curve-billed Thrasher, prolonged periods of acute stress may be limited to the Sonoran Desert portion of their range. Among these three, the wren and thrasher are insectivores and so have constant access to preformed water in their diet, whereas the granivorous Abert's Towhee may have very limited access to water resources.

**Other Factors Affecting Species Distributions in Hot Deserts.** Avoidance of environmental extremes is a common strategy used by animals to balance their water and energy budgets under physiologically challenging conditions, and the availability of thermally buffered microsites is potentially important in modeling the consequences of hotter future climates. During the hottest part of the day, small birds completely cease all foraging activities (18, 19) and seek relatively cool, shaded thermal refugia to reduce exposure to extreme heat (20, 21). Our model focuses on meteorological  $T_{air}$ , which for birds in hot desert environments, is typically a minimum index of thermal stress in the absence of radiative and convective heat fluxes, and approximates the thermal conditions in a completely shaded and ventilated meteorological thermoscreen. There is also evidence that North American Land Data Assimilation System (NLDAS) and other gridded climate products often underestimate maximum temperatures by  $1^\circ\text{C}$  or more (22), underscoring the conservative nature of our analysis. Nevertheless, it may be possible for individuals to use cooler microsites (e.g., burrows and sites in dense vegetation) (23). The role of microsites is emphasized by the observation that, in hot areas like the Sonoran Desert, many species already experience periods of significant dehydration risk on a small number of days per year (Fig. 2). However, the availability of microsites, given habitat conditions and competition, may not be guaranteed, presently or under future environmental changes.

The increased mortality risk associated with acute heat stress under climate change that we have modeled here will be superimposed on a range of fitness impacts of chronic, but sublethal on a daily scale, heat exposure that we are only just beginning to understand; these include reductions in body condition driven by tradeoffs between foraging efficiency and heat dissipation behaviors (24), and temperature effects on offspring provisioning rates with consequences for chick development and survival (25). Evidence from arid and semiarid ecosystems indicates that shifts in avian abundance associated with heat waves and drought are already occurring (1, 26, 27).

In this study, we have focused on arid environments and modeled EWL under conditions near the ceiling of avian physiological capacity. However, the increases in the frequency dehydration risk for Lesser Goldfinch in the humid eastern half of the United States (Fig. S1) raises the question of how humidity will modify the impacts of higher air temperatures. By reducing the efficiency of evaporative cooling during extremely hot conditions (28), elevated humidity will increase the likelihood that birds cannot dissipate heat rapidly

**Table 1. Sampled passerine species and measured body masses**

Name	Mass (g; $\bar{X} \pm \text{SD}$ )	N	Notes ( <a href="https://www.allaboutbirds.org">https://www.allaboutbirds.org</a> )
Lesser Goldfinch ( <i>Spinus psaltria</i> )	9.7 $\pm$ 0.5	33	Open habitat dweller, granivore, tree-nester, foliage gleaner
House Finch ( <i>Haemorhous mexicanus</i> )	18.0 $\pm$ 1.3	75	Human/wild habitat dweller, granivore, tree-nesting, ground forager
Cactus Wren ( <i>Campylorhynchus brunneicapillus</i> )	34.6 $\pm$ 2.7	22	Desert dweller, insectivore, shrub-nesting, ground forager
Abert's Towhee ( <i>Melospiza aberti</i> )	41.8 $\pm$ 1.3	28	Desert riparian habitat dweller, granivore, tree-nester, ground forager
Curve-billed Thrasher ( <i>Toxostoma curvirostre</i> )	71.0 $\pm$ 8	18	Scrub habitat dweller, insectivore, shrub-nester, ground forager

enough to avoid lethal hyperthermia. Extremely hot, humid weather will create conditions in which avian mortality events analogous to those that have affected pteropodid bats along the east coast of Australia (29) may be expected to become more frequent.

**Future Changes in Avian Distributions and Community Structure.** The 4 °C warmer climate scenario produced dramatic increases in the frequency, intensity, and extent of lethal dehydration conditions. These conditions may overwhelm population-level (e.g., density dependence and metapopulation dynamics) and individual-level (behavior and movement) coping mechanisms. The spatial expansion of lethal dehydration may be of greatest concern for conservation of desert avifaunas. At an individual level, birds may be able to cope with extreme heat events near the lower end of the severity range we have modeled here. At a population/species level, birds may be able to cope with more severe scenarios (e.g.,  $\leq 3$ -h survival times), because such extreme events are unlikely to affect entire ranges simultaneously. However, when such intense events begin to occur on a regular and sustained basis, as will be the case for Lesser Goldfinch and other small passerines later in this century, one must question the ability of birds smaller than 10–20 g to avoid extirpation in some portions of their US Southwest range. The relationship between body size and dehydration risk may ultimately drive shifts in community structure in hot regions, as smaller birds experience more rapid declines in survival and productivity compared with larger species.

**Conservation Implications.** These findings suggest several key conservation and management implications for avifaunas of the desert Southwest and other hot desert regions of the world. The US Southwest enjoys a relatively high avian diversity, distinctive avian assemblages, and a number of threatened birds that may be vulnerable to extreme heat waves (30). The present model was limited to the US portion of the range of our focal species. However, we expect that the more southern Mexican portions of Sonoran and Chihuahuan species' ranges may experience similar current and future regimes of high temperature extremes. In general, because passerine birds rely on the metabolically costly process of panting to evaporate water and lack more efficient evaporative mechanisms (e.g., gular flutter and cutaneous evaporation) (12, 31), small species may experience a new level of threat from hot extremes in this region.

This study provides further motivation for limiting the magnitude of climate warming, because there is already evidence the majority of high temperature extremes are attributable to anthropogenic warming (32). Our study also identifies an urgent need for conservation professionals to consider the effects of hot extremes on desert fauna and promote adaptive measures. For example, efforts to conserve vegetation providing thermal refugia and the maintenance of sources of open water may be important. Similarly, we emphasize that conservation planning should consider the implications of high temperature extremes and climate change by ensuring the availability and conservation of more thermally moderate habitats.

## Materials and Methods

**Evaporative Water Loss Measurements.** We conducted evaporative water loss experiments on 193 individuals of five passerine bird species during June and July of 2012 and 2013 in an area just north of Tucson, AZ (Table 1;  $\sim 32^\circ 30' \text{N}$ ,

$110^\circ 55' \text{W}$ ). These regionally common species range from 9.7 to 71.0 g and include insectivore and granivore species. During the sampling period at the Tucson, AZ, MET weather station, daily maximum  $T_{\text{air}}$  ranged from 31 to 42 °C, and the dew point ranged from  $-15$  to 23 °C.

Experiments were conducted with the approval of University of New Mexico Institutional Animal Care and Use Committee (protocol no. 12-100537-MCC) and under permits issued by the US Fish and Wildlife Service and the Arizona Game and Fish Department. The birds were captured using standard walk-in traps baited with seed; upon capture, the birds were held outdoors in shaded, darkened screen cages. All birds were subjected to thermoregulatory trials and released on the same day of capture.

Before each experimental run, a temperature-sensitive passive integrated transponder (PIT) tag (model TX1411BBT; Biomark) was injected into each bird's abdominal cavity and used to monitor  $T_{\text{b}}$  during trials (33). Measurements of resting metabolic rate and EWL were made using a flow-through respirometry system (33). The respirometry chamber consisted of a transparent plastic container (5 L) sealed by an opaque lid and modified by addition of inlet and outlet ports; it contained a stainless steel hardware cloth floor above a 2-cm layer of medium-weight mineral oil to trap excreta. The respirometry chamber was placed inside a modified ice chest where a microprocessor-controlled Peltier device regulated  $T_{\text{air}}$  to  $\pm 0.5$  °C. Dry air provided by a compressor and pushed through a membrane air dryer was supplied to a mass flow controller that regulated airflow to the respirometry chamber at a rate that ranged from 5 to 40 standard liters per minute. Chamber excurrent chamber air was sampled at 250 mL·min<sup>-1</sup> and directed to a CO<sub>2</sub>/H<sub>2</sub>O gas analyzer (model LI-840A; LICOR) with an accuracy of  $\pm 1$  ppm for CO<sub>2</sub> and 1% of the reading for H<sub>2</sub>O. Rates of whole-animal EWL (grams of water per hour) were calculated from the H<sub>2</sub>O ppt readings (less baseline values) for the same data points using equation 10.9 in Lighton (34).

Following PIT tag insertion, each bird was weighed ( $\pm 0.01$  g) and placed in the respirometry chamber. The chamber was darkened during runs and an infrared light and video camera enabled continuous viewing of the animal in the chamber on a computer monitor. The bird was exposed to one or more thermoneutral temperatures (30 °C or 35 °C) and one or more higher temperatures (40–54 °C in 2 °C increments) in a continuously stepped pattern of temperature trials over the span of 1–3 h. To maintain H<sub>2</sub>O content of the respirometry chamber at values that would not impede evaporation (dew point  $< 5$  °C), flow rates were maintained between 5 and 40 L·min<sup>-1</sup>, depending on the chamber temperature and the bird's evaporative rate. The initial thermoneutral trial temperature allowed a bird to calm from handling; H<sub>2</sub>O and CO<sub>2</sub> production were monitored and observed to drop to resting levels (typically 30–45 min). Temperatures were then increased and birds were allowed to adjust to each for 10–20 min and stabilize before moving to the next temperature. Most birds had few problems coping with  $T_{\text{air}}$  below 46 °C, but at higher temperatures, birds sometimes showed intermittent activity such as flapping escape attempts; these periods were not included in the analysis. Because airflow rates through the chamber ranged from 5 to 40 L·min<sup>-1</sup>, chamber equilibrium times ranged from  $\sim 30$  s to 4.5 min (35), which allowed for accurate observations of both activity metabolism and resting metabolism. As chamber  $T_{\text{air}}$  were increased above thermoneutrality,  $T_{\text{b}}$  and activity were monitored closely to prevent mortality. Experimental runs were also timed to end before total losses, including fecal, were less than 10% of  $M_{\text{b}}$  to minimize the effects of hypohydration on thermoregulatory performance.

**Geospatial Temperature Time Series.** To characterize 1979–2012  $T_{\text{air}}$  across the conterminous US study region, we used the multiinstitution NLDAS-2 (36). In this dataset, the temperature forcing fields are spatially interpolated from 32-km resolution to a 1/8° grid ( $\sim 14$  km) and temporally disaggregated from 3-h to hourly frequency. This combination of spatial and temporal resolution makes NLDAS-2 well suited for simulation of many ecologically relevant land surface processes over large geographic areas. The data are available from 1979 onward on the NASA Goddard Earth Science Data and Information Services Center website (<https://disc.sci.gsfc.nasa.gov/>).

**Water Balance Model.** At  $T_{\text{air}}$  below normothermic  $T_b$ , EWL rates are typically modest and determined by the obligatory losses associated with gas exchange at respiratory surfaces and minimum evaporative losses across the skin. Most heat loss is passive via conduction, convection, and radiation, and determined by the thermal gradient between the animal and its environment. As environmental temperatures exceed  $T_b$ , rates of EWL increase rapidly because both metabolic and environmental heat loads must be dissipated solely through evaporative processes. In passerines, this active heat dissipation is characterized by panting, which increases both EWL and metabolic rate. As  $T_{\text{air}}$  exceeds  $T_b$ , birds retreat to shaded microsites and often remain inactive during the heat of the day; this minimizes EWL, but water intake from foraging is also curtailed. In this model, we assume that birds remain in shaded areas during the hottest periods of the day and forgo foraging. For simplicity, we use  $T_{\text{air}}$  to define the thermal stress on the animal and its rate of evaporative water loss (12, 37), while recognizing that complex thermal and radiative environments outside of still-shaded microsites may produce additional thermal stress.

For each species, a linear regression model was fitted to EWL rate against chamber  $T_{\text{air}} > \sim 40^\circ\text{C}$  where active evaporative heat dissipation is occurring. The resulting linear model for each species was used to estimate hourly EWL rates and water balance at  $T_{\text{air}}$  above upper critical temperature ( $T_{\text{uc}}$ ). In this analysis, we estimate the total evaporative water loss by the animal between 10 AM and 6 PM (8 h) local time, a period when continued foraging would likely produce a net water deficit because of increased heat loads during high-temperature events. We view this estimate as conservative because  $T_{\text{air}}$  in this region may exceed  $40^\circ\text{C}$  for 10 or more hours each day during a heat wave.

**Survival Time Modeling.** We use the concept of survival time as an indicator of the intensity of a high-temperature event and the probability of death. Survival time was estimated as the number of hours during which cumulative water loss via evaporation was equivalent to  $<15\%$   $M_b$  for each passerine species. Given daily activity patterns and temperature profiles, we focus on

two ecologically important survival times:  $\leq 5$  h (moderate risk of lethal dehydration) and  $\leq 3$  h (high risk). Although data for birds are sparse, we selected the loss of  $15\%$  of  $M_b$  as a conservative estimate of dehydration tolerance (21, 38), and assume that mortality occurs when this threshold is exceeded. For each species, survival times during warm season (April to September) were calculated daily at each NLDAS-2 grid cell location. Our measurements of EWL subsume the effects of hyperthermia, but we do not assess the chronic effects heat on potential mortality. Interactive Data Language code is available in [Dataset S1](#).

**Regional and Climate-Change Analyses.** With the goal of identifying relevant areas prone to lethal dehydration, we obtained bird ranges from US Geological Survey National Gap Analysis Program portal (<https://gapanalysis.usgs.gov/species/data/download/>) and overlaid these with the NLDAS-2 grid. We calculated and mapped regional summaries (dehydration climatologies) across species' US ranges as well as within the more-restricted southwest desert ecoregion comprising the Mojave, Colorado, Sonoran, and Chihuahuan deserts (39) to minimize the influence of variations in range size across our suite of species. We then repeated dehydration and mapping analyses using a warming scenario, wherein all NLDAS-2 temperatures during our 1979–2012 dataset were increased by  $4^\circ\text{C}$ , in line with current regional projections for summer warming by 2070–2099 under Special Report on Emissions Scenarios A2 (14) and with mean global surface temperature increases under Representative Concentration Pathway 8.5 (9).

**ACKNOWLEDGMENTS.** We thank G. Sadoti and S. A. McAfee for helpful discussions and the Cadden and Wolf families for logistical support and access to their land for research. This work was supported by National Science Foundation Grant IOS-1122228 (to B.O.W.) and National Aeronautics and Space Administration Grant NNX13AB65G (to T.P.A.). A.R.G. was supported by Natural Sciences and Engineering Research Council of Canada Postdoctoral Fellowship 420761.

- Albright TP, et al. (2010) Combined effects of heat waves and droughts on avian communities across the conterminous United States. *Ecosphere*, 10.1890/ES10-00057.1.
- Jentsch A, Beierkuhnlein C (2008) Research frontiers in climate change: Effects of extreme meteorological events on ecosystems. *C R Geosci* 340(9-10):621–628.
- Ma G, Rudolf VHW, Ma C-S (2014) Extreme temperature events alter demographic rates, relative fitness, and community structure. *Glob Chang Biol* 21:1794–1808.
- Coumou D, Robinson A (2013) Historic and future increase in the global land area affected by monthly heat extremes. *Environ Res Lett* 8(3):034018.
- Mutibwa D, Vavrus SJ, McAfee SA, Albright TP (2015) Recent spatiotemporal patterns in temperature extremes across conterminous United States. *J Geophys Res Atmos* 120(15):7378–7392.
- McKechnie AE, Hockey PAR, Wolf BO (2012) Feeling the heat: Australian landbirds and climate change. *Emu* 112(2):i–vii.
- Fey SB, et al. (2015) Recent shifts in the occurrence, cause, and magnitude of animal mass mortality events. *Proc Natl Acad Sci USA* 112(4):1083–1088.
- Coumou D, Rahmstorf S (2012) A decade of weather extremes. *Nat Clim Chang* 2(7):491–496.
- Intergovernmental Panel on Climate Change (IPCC) (2013) Summary for Policymakers. *Climate Change 2013: The Physical Science Basis*, Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, eds Stocker TF, et al. (Cambridge Univ Press, Cambridge, UK).
- McKechnie AE, Wolf BO (2010) Climate change increases the likelihood of catastrophic avian mortality events during extreme heat waves. *Biol Lett* 6(2):253–256.
- Calder WA, King JR (1974) Thermal and caloric relations of birds. *Avian Biology*, ed Farner DS, King JR (Academic, New York), Vol 4, pp 259–413.
- Wolf B, Walsberg G (1996) Respiratory and cutaneous evaporative water loss at high environmental temperatures in a small bird. *J Exp Biol* 199(Pt 2):451–457.
- Wolf BO, Walsberg GE (1996) Thermal effects of radiation and wind on a small bird and implications for microsite selection. *Ecology* 77(7):2228–2236.
- Walsh J, et al. (2014) Our Changing Climate. Climate Change Impacts in the United States: The Third National Climate Assessment. Available at [www.globalchange.gov/browse/reports/climate-change-impacts-united-states-third-national-climate-assessment-0](http://www.globalchange.gov/browse/reports/climate-change-impacts-united-states-third-national-climate-assessment-0). Accessed May 16, 2014.
- Dawson WR (1982) Evaporative losses of water by birds. *Comp Biochem Physiol A* 71(4):495–509.
- Williams JB (1996) A phylogenetic perspective of evaporative water loss in birds. *Auk* 113(2):457–472.
- Whitfield MC, Smit B, McKechnie AE, Wolf BO (2015) Avian thermoregulation in the heat: Scaling of heat tolerance and evaporative cooling capacity in three southern African arid-zone passerines. *J Exp Biol* 218(Pt 11):1705–1714.
- Ricklefs RE, Hainsworth FR (1969) Temperature regulation in nestling cactus wrens: The nest environment. *Condor* 71(1):32–37.
- Austin G (1978) Daily time budget of the postnesting Verdin. *Auk* 95(2):247–251.
- Wolf BO, Wooden KM, Walsberg GE (1996) The use of thermal refugia by two small desert birds. *Condor* 98(2):424–428.
- Wolf B (2000) Global warming and avian occupancy of hot deserts: A physiological and behavioral perspective. *Rev Chil Hist Nat* 73:395–400.
- Behnke R, et al. (2016) Evaluation of downscaled, gridded climate data for the conterminous United States. *Ecol Appl* 26(5):1338–1351.
- Scheffers BR, Edwards DP, Diesmos A, Williams SE, Evans TA (2014) Microhabitats reduce animal's exposure to climate extremes. *Glob Chang Biol* 20(2):495–503.
- du Plessis KL, Martin RO, Hockey PAR, Cunningham SJ, Ridley AR (2012) The costs of keeping cool in a warming world: Implications of high temperatures for foraging, thermoregulation and body condition of an arid-zone bird. *Glob Chang Biol* 18(10):3063–3070.
- Cunningham SJ, Kruger AC, Nxumalo MP, Hockey PA (2013) Identifying biologically meaningful hot-weather events using threshold temperatures that affect life-history. *PLoS One* 8(12):e82492.
- George TL, Fowler AC, Knight RL, McEwen LC (1992) Impacts of a severe drought on grassland birds in western North Dakota. *Ecol Appl* 2(3):275–284.
- Cruz-MacDonnell KK, Wolf BO (2015) Rapid warming and drought negatively impact population size and reproductive dynamics of an avian predator in the arid southwest. *Glob Chang Biol* 22:237–253.
- Gerson AR, Smith EK, Smit B, McKechnie AE, Wolf BO (2014) The impact of humidity on evaporative cooling in small desert birds exposed to high air temperatures. *Physiol Biochem Zool* 87(6):782–795.
- Welbergen JA, Klose SM, Markus N, Eby P (2008) Climate change and the effects of temperature extremes on Australian flying-foxes. *Proc Biol Sci* 275(1633):419–425.
- Jenkins CN, Van Houtan KS, Pimm SL, Sexton JO (2015) US protected lands mismatch biodiversity priorities. *Proc Natl Acad Sci USA* 112(16):5081–5086.
- Tieleman BI, Williams JB (2002) Cutaneous and respiratory water loss in larks from arid and mesic environments. *Physiol Biochem Zool* 75(6):590–599.
- Fischer EM, Knutti R (2015) Anthropogenic contribution to global occurrence of heavy-precipitation and high-temperature extremes. *Nat Clim Chang* 5:560–564.
- Smith EK, O'Neill J, Gerson AR, Wolf BO (2015) Avian thermoregulation in the heat: Resting metabolism, evaporative cooling and heat tolerance in Sonoran Desert doves and quail. *J Exp Biol* 218(Pt 22):3636–3646.
- Lighton J (2008) *Measuring Metabolic Rates* (Oxford Univ Press, Oxford, UK).
- Lasiewski RC, Acosta AL, Bernstein MH (1966) Evaporative water loss in birds—I. Characteristics of the open flow method of determination, and their relation to estimates of thermoregulatory ability. *Comp Biochem Physiol* 19(2):445–457.
- Cosgrove BA, et al. (2003) Real-time and retrospective forcing in the North American Land Data Assimilation System (NLDAS) project. *J Geophys Res* 108(D22):8842.
- Ricklefs RE, Hainsworth FR (1968) Temperature regulation in Nestling Cactus Wrens: The development of homeothermy. *Condor* 70(2):121–127.
- Arad Z, Horowitz M, Eylath U, Marder J (1989) Osmoregulation and body fluid compartmentalization in dehydrated heat-exposed pigeons. *Am J Physiol* 257(2 Pt 2):R377–R382.
- Bailey RG (1980) *Description of the Ecoregions of the United States* (USDA Forest Service, Washington, DC).